

# Crixivan



Crixivan is used to help treat HIV infection by blocking the action of protease, an enzyme which the virus needs to multiply.

Prices may vary slightly when paying.

Crixivan 400 mg x 10 pill **SGD92.11** [Buy now](#)

Crixivan 400 mg x 30 pill **SGD227.42** [Buy now](#)

Crixivan 400 mg x 60 pill **SGD435.3** [Buy now](#)

## Description

### General information

Crixivan goes to some type of drugs referred to as protease inhibitors. It's used in conjunction with other Aids medicines to deal with the problem triggered through the hiv (Aids). Aids may be the virus accountable for acquired immune deficiency syndrome (Helps). Crixivan isn't a remedy for Aids. It doesn't avoid the spread of Aids to other people through sexual contact or bloodstream contamination. Medicines could also be used to avoid Aids infection after connection with herpes.

### Directions

Take Crixivan just as recommended from your physician. Take medicines orally having a full glass water. The suggested adult dose is 800 mg taken with water every 8 hrs. In individuals with mild to moderate liver disease, the suggested dose is 600 mg every eight hrs. It ought to be taken 1 hour before or 2 hrs following a meal. To assist decrease the chance of kidney gemstones, drink a minimum of 6 full portions of water or any other fluids during the day.

### Safeguards

Before you take Crixivan you need to engage with your physician for those who have kidney gemstones, liver disease, liver failure, cirrhosis, hepatitis, diabetes, hemophilia, diabetes, heart disease, coronary heart, cardiac arrest, kidney disease, kidney failure, any allergic reactions. Children might be more responsive to the unwanted effects of the drug. Taking indinavir won't stop you from passing Aids with other people through unguaranteed sex or discussing of needles. Avoid using this medication having a meal full of calories, body fat, and protein because this may decrease its effect.

### Advisable limitations

Don't take Crixivan if you're allergic to indinavir or any elements of the medication, take the following medicines: alprazolam, amiodarone, astemizole, cisapride, ergot types (ergotamine, dihydroergotamine), midazolam, pimozone, terfenadine, triazolam. Enable your physician know if you're pregnant or breastfeeding.

### Possible side-effect

Get emergency medical help for those who have stomach discomfort, nausea, vomiting, acid reflux, appetite loss, headache, lightheadedness, drowsiness, weakness, confusion, lack of fluids, mouth sores, muscle aches or discomfort, pounding heartbeat, elevated thirst, elevated peeing, yellowing of your skin or eyes, weight reduction, painful peeing, bloodstream in urine, dark urine, pale stools, chest discomfort or pressure, peeling, blistering, back discomfort, chills, fever, rash, itchiness, swelling from the face, tongue, throat, trouble breathing. If you see other effects unlisted above, speak to your physician.

### Drug interactions

Inform your physician about other medicines you utilize, especially: seizure medicines (phenytoin, felbamate, carbamazepine), Aids/Helps medicines (nevirapine, lamivudine, ritonavir, zidovudine), phosphodiesterase type-5 inhibitors (tadalafil, sildenafil, vardenafil), blood insulin, heart rhythm medicines (quinidine, flecainide, amiodarone), cyclosporine, tacrolimus, cholesterol-lowering medications (fluvastatin, cerivastatin, simvastatin, lovastatin), calcium funnel blockers (nimodipine, verapamil, nifedipine), mao inhibitors (fluoxetine,

amitriptyline, citalopram, nortriptyline, doxepin), anti-biotics (telithromycin, erythromycin, clarithromycin), fluticasone. Interaction between two medicines doesn't imply that you need to quit taking one of these. Inform your physician or prescriber about all prescription, over-the-counter, and herbal medicines you're taking.

#### Skipped dose

Go ahead and take skipped dose as quickly as possible. Skip the skipped dose if it's time for your forthcoming scheduled dose. Do not take extra medicine to from the skipped dose.

#### Overdose

If you feel you've overdosed the medication seek emergency medical help at the same time. The overdose signs and symptoms are nausea, vomiting, diarrhea, back discomfort, bloodstream within the urine.

#### Storage

Keep medicine at 70 degrees between 59-86 levels F (15-30 levels C) from light and moisture. Don't keep drugs within the bathroom. Keep all drugs from achieve of kids and pets.

#### Note

The data presented to begin includes a general character. Note please these details can't be employed for self-treatment and self diagnosis. You need to talk to your physician or healthcare agent regarding any sort of instructions of the condition. The details are reliable, but we concede it might contain mistakes. We're not accountable for any direct, indirect, special or any other damage triggered by utilization of these details on the website and for effects of self-treatment.

#### Tags

[buy Crixivan in chinatown](#), [Crixivan pills](#), [Crixivan online singapore](#), [buy Crixivan in tanglin](#), [buy Crixivan in Jurong](#), [buy Crixivan in Bukit Merah](#), [buy Crixivan in Hougang](#), [cheap Crixivan online Singapore](#), [Crixivan fast delivery](#), [Crixivan altrnatives](#), [buy Crixivan in Woodlands](#), [buy Crixivan in Kuala Lumpur](#), [Crixivan online paypal](#), [buy Crixivan online without prescription](#), [Crixivan tablets online Singapore](#), [buy Crixivan online in Singapore](#), [buy Crixivan online mastercard](#), [buy Crixivan overnight](#), [buy Crixivan in keppel](#), [buy Crixivan 400 mg in Singapore](#), [Crixivan online pharmacy](#), [order Crixivan online in Singapore](#), [Crixivan for sale](#), [cheapest Crixivan online Singapore](#), [buy Crixivan in Bedok](#), [buy Crixivan online with no prescription](#), [generic Crixivan online](#),