

Detrol La



Detrol La is an antispasmodic that is used to treat an overactive bladder by relaxing the muscles in the bladder.

Prices may vary slightly when paying.

Detrol La 4 mg x 10 pill	SGD69.47 Buy now
Detrol La 4 mg x 30 pill	SGD160.32 Buy now
Detrol La 4 mg x 60 pill	SGD287.11 Buy now
Detrol La 4 mg x 90 pill	SGD408.31 Buy now
Detrol La 4 mg x 120 pill	SGD540.14 Buy now

Description

General information

Detrol La goes to several medicines referred to as antispasmodics and anticholinergics. Medicines can be used to deal with an over active bladder. By relaxing the muscles within the bladder, tolterodine enhances the opportunity to control the peeing. It cuts down on spasms from the bladder muscles.

Directions

Take Detrol La just as recommended from your physician. The typical suggested dose is 4 mg once daily to begin. The dose may later be reduced to two mg once daily according to individual response and requires. The utmost dose is 4 mg daily. For those who have reduced liver or kidney function, and individuals taking certain medicines, the suggested dose is 2 mg once daily. Take medicines orally, without or with food. Go having a full glass water. Swallow the pill whole.

Safeguards

Before you take Detrol La you need to engage with your physician for those who have kidney disease, kidney failure, liver disease, liver failure, cirrhosis, hepatitis, irregular heart rhythm (arrhythmia), glaucoma, Myasthenia gravis, bloating, difficulty urinating, enlarged prostate (benign prostatic hyperplasia), any allergic reactions. It may cause blurred vision or unwanted effects that could impair your opinions or responses. Don't drive or perform tasks that need performance. Limit alcoholic drinks. Alcohol can increase a few of the unwanted effects. Medicines may cause decreased sweating. Seniors people might be more responsive to this drug.

Advisable limitations

You shouldn't take Detrol La if you're allergic to tolterodine or the elements from the medication, or you have gastric retention (stomach obstruction), out of control narrow-position glaucoma, urinary retention. Enable your physician know if you're pregnant or breastfeeding.

Possible side-effect

Get emergency medical help for those who have xerostomia, dry eyes, headache, constipation, stomach upset or discomfort, lightheadedness, sleepiness, fatigue, blurred vision, fatigue, gas, abdominal discomfort, acid reflux, bloody urine, trouble peeing, chest discomfort, irregular heartbeat, fainting, swelling from the face or throat, hives, breathlessness, fever, itchiness. If you see other effects unlisted above, speak to your physician.

Drug interactions

Inform your physician about other medicines you utilize, especially: heart rhythm medicine (procainamide, sotalol, quinidine, dofetilide), antipsychotic medicines (chlorpromazine, thiothixene, loxapine), anti-biotics (clarithromycin, telithromycin, erythromycin), vinblastine, pentamidine, narcotic medicines (methadone, levomethadyl), droperidol, cyclosporine, chloroquine, halofantrine, anticholinergic medicines (clidinium, atropine, dicyclomine, belladonna, scopolamine), protease inhibitors (atazanavir, indinavir, ritonavir,

amprenavir). Interaction between two medicines doesn't imply that you need to quit taking one of these. Inform your physician or prescriber about all prescription, over-the-counter, and herbal medicines you're taking.

Skipped dose

Go ahead and take skipped dose as quickly as possible. Skip the skipped dose if it's time for your forthcoming scheduled dose. Do not take extra medicine to from the skipped dose.

Overdose

If you feel you've overdosed the medication seek emergency medical help at the same time. The overdose signs and symptoms are fast, slow or irregular heartbeat, fainting, unusual excitement, mental or mood changes.

Storage

Keep medicine at 70 degrees 77 levels F (25 levels C) from light and moisture. Don't keep drugs within the bathroom. Keep all drugs from achieve of kids and pets.

Note

The data presented to begin includes a general character. Note please these details can't be employed for self-treatment and self diagnosis. You need to talk to your physician or healthcare agent regarding any sort of instructions of the condition. The details are reliable, but we concede it might contain mistakes. We're not accountable for any direct, indirect, special or any other damage triggered by utilization of these details on the website and for effects of self-treatment.

Tags

[buy Detrol La 4 mg x 30 pill in Singapore](#), [buy Detrol La online without prescription](#), [buy Detrol La in Bedok](#), [buy Detrol La in Woodlands](#), [buy Detrol La in Kuala Lumpur](#), [Detrol La online singapore](#), [buy Detrol La 4 mg x 120 pill in Singapore](#), [buy Detrol La in Hougang](#), [Detrol La online pharmacy](#), [buy Detrol La in keppel](#), [Detrol La pills](#), [buy Detrol La in changi](#), [buy Detrol La online with no prescription](#), [Detrol La fast delivery](#), [buy Detrol La in tanglin](#), [buy Detrol La online in Singapore](#), [generic Detrol La online](#), [buy Detrol La 4 mg in Singapore](#), [buy Detrol La overnight](#), [buy Detrol La online mastercard](#), [buy Detrol La in chu kang](#), [Detrol La tablets online Singapore](#), [buy Detrol La in Bukit Merah](#), [Detrol La altrnatives](#), [buy Detrol La in toa payoh](#), [order Detrol La online in Singapore](#), [Detrol La for sale](#), [Detrol La online paypal](#), [cheap Detrol La online Singapore](#), [buy Detrol La in Jurong](#), [cheapest Detrol La online Singapore](#),