

Indinavir



Indinavir is an antiviral medication used to treat HIV, which causes AIDS by preventing HIV cells from multiplying in your body.

Prices may vary slightly when paying.

Indinavir 400 mg x 30 pill **SGD136** [Buy now](#)

Indinavir 400 mg x 60 pill **SGD206.52** [Buy now](#)

Description

General information

Indinavir goes towards the type of medicines known as protease inhibitors. It's used in conjunction with other Aids medicines to deal with the problem triggered through the hiv (Aids). Aids may be the virus accountable for acquired immune deficiency syndrome (Helps). Additionally, it reduces your chance of getting Aids complications (for example new infections, cancer). Medicines could also be used to avoid Aids infection after connection with herpes. Indinavir isn't a remedy for Aids. It doesn't avoid the spread of Aids to other people through sexual contact or bloodstream contamination.

Directions

Take Indinavir just as recommended from your physician. Take medicines orally having a full glass water before eating anything (a minimum of one hour before or 2 hrs after food), usually 3 occasions daily. Drink a minimum of 6 full portions of water or any other fluids during the day to lower the chance of kidney gemstones. The suggested indinavir dose for dealing with grown ups with Aids or Helps is 800 mg (two 400 mg capsules) every eight hrs. If you're taking certain other medicines, a lesser dose might be suggested. The suggested dose for those who have liver disease is 600 mg every eight hrs.

Safeguards

Before you take Indinavir you need to engage with your physician for those who have kidney gemstones, liver disease, liver failure, cirrhosis, hepatitis, diabetes, hemophilia, coronary heart, cardiac arrest, kidney disease, kidney failure, high bloodstream body fat levels, any allergic reactions. Taking indinavir won't stop you from passing Aids with other people through unguaranteed sex or discussing of needles. Children might be more responsive to the unwanted effects of the drug.

Advisable limitations

Don't take Indinavir if you're allergic to indinavir or any elements of the medication or take the following medicines: alprazolam, amiodarone, astemizole, cisapride, ergot types, midazolam, pimozide, terfenadine, triazolam. Enable your physician know if you're pregnant or breastfeeding.

Possible side-effect

Get emergency medical help for those who have dry or scratchy skin, abdominal or stomach discomfort, difficulty in sleeping, acidity reflux, nausea, acid reflux, diarrhea, appetite loss, bloody urine, elevated peeing, hives, breathlessness, swelling of the face, lips, tongue, constipation, lightheadedness, weakness, headache, drowsiness, rash, vomiting, jaundice (yellowing of your skin or eyes), easy bruising, mood changes, tired feeling, bloatedness, numbness, weight reduction, pale stools, peeling, back discomfort, chills, fever, rapid heartbeat, inflamed abdomen. If you see other effects unlisted above, speak to your physician.

Drug interactions

Inform your physician about other medicines you utilize, especially: seizure medicines (phenytoin, felbamate, carbamazepine), Aids/Helps medicines (nevirapine, lamivudine, zidovudine), drugs for male erectile problems (tadalafil, vardenafil, sildenafil), blood insulin, heart rhythm medicines (quinidine, disopyramide, amiodarone), cyclosporine, tacrolimus, cholesterol-lowering medications (fluvastatin, cerivastatin, simvastatin, lovastatin), calcium funnel blockers (nimodipine, verapamil, nifedipine), mao inhibitors (fluoxetine, amitriptyline,

citalopram, nortriptyline, doxepin), anti-biotics (telithromycin, erythromycin, clarithromycin), antifungal medicines (fluconazole, ketoconazole, itraconazole), fluticasone. Interaction between two medicines doesn't imply that you need to quit taking one of these. Inform your physician or prescriber about all prescription, over-the-counter, and herbal medicines you're taking.

Skipped dose

Go ahead and take skipped dose as quickly as possible. Skip the skipped dose if it's time for your forthcoming scheduled dose. Do not take extra medicine to from the skipped dose.

Overdose

If you feel you've overdosed the medication seek emergency medical help at the same time. The overdose signs and symptoms are nausea, emesis, diarrhea, back pain, bloody urine.

Storage

Keep medicine at 70 degrees between 59-77 levels F (15-30 levels C) from light and moisture. Don't keep drugs within the bathroom. Keep all drugs from achieve of kids and pets.

Note

The data presented to begin includes a general character. Note please these details can't be employed for self-treatment and self diagnosis. You need to talk to your physician or healthcare agent regarding any sort of instructions of the condition. The details are reliable, but we concede it might contain mistakes. We're not accountable for any direct, indirect, special or any other damage triggered by utilization of these details on the website and for effects of self-treatment.

Tags

[Indinavir for sale](#), [buy Indinavir in cbd](#), [order Indinavir online in Singapore](#), [Indinavir altrnatives](#), [buy Indinavir online mastercard](#), [buy Indinavir in Bedok](#), [buy Indinavir in chinatown](#), [cheapest Indinavir online Singapore](#), [buy Indinavir in Woodlands](#), [buy Indinavir in marina area](#), [Indinavir online singapore](#), [generic Indinavir online](#), [buy Indinavir in chu kang](#), [buy Indinavir in Hougang](#), [buy Indinavir overnight](#), [cheap Indinavir online Singapore](#), [buy Indinavir online with no prescription](#), [buy Indinavir online without prescription](#), [buy Indinavir 400 mg in Singapore](#), [Indinavir pills](#), [buy Indinavir online in Singapore](#), [buy Indinavir in Kuala Lumpur](#), [Indinavir fast delivery](#), [Indinavir online pharmacy](#), [Indinavir online paypal](#), [buy Indinavir in changi](#), [buy Indinavir in keppel](#), [Indinavir tablets online Singapore](#), [buy Indinavir in Bukit Merah](#), [buy Indinavir in Jurong](#),